Universe International Journal of Interdisciplinary Research
(International Peer Reviewed Refereed Journal)
DOI No. – 08.2020-25662434

© UIJIR | ISSN (O) – 2582-6417
MARCH 2024 | Vol. 4 Issue 10
www.uijir.com

INVESTIGATING THE IMPACT OF ANTIBIOTICS ON CHILD IMMUNITY: UNRAVELING MECHANISMS AND IMPLICATIONS FOR HEALTH

Author's Name: Bhumi Pandya [1]

Affiliation:

1. Principal, Shankersinh Vaghela Bapu Institute of Nursing, Gandhinagar, Gujarat, India.

Corresponding Author Name and Email Id: Bhumi Pandya,
bhumipandya46@yahoo.com

ABSTRACT

The widespread utilization of antibiotics in pediatric populations has emerged as a subject of concern due to its potential impact on child immunity. This comprehensive review aims to consolidate current understanding surrounding the effects of antibiotics on the developing immune system in children, delving into the underlying mechanisms and evaluating implications for health outcomes. Central to this exploration is the elucidation of how antibiotics perturb microbiota composition, disrupt immune homeostasis, and influence immune maturation during critical developmental stages. Moreover, the review scrutinizes the long-term ramifications of early-life antibiotic exposure on immune resilience, susceptibility to infections, and the development of immune-related disorders in later life. Additionally, the examination extends to potential strategies aimed at mitigating the adverse effects of antibiotics on child immunity, while concurrently preserving their therapeutic benefits against bacterial infections. By highlighting the necessity of optimizing antibiotic use in children, this review underscores the imperative to safeguard immune health and promote long-term well-being in pediatric populations.

Keywords: : Antibiotics, Child immunity, Health implications, Immune maturation, Immune resilience, Susceptibility to infections, Immune-related disorders, Therapeutic benefits, Optimization of antibiotic use.

DOI Link :: https://doi-ds.org/doilink/03.2024-51788784/UIJIR

© UIJIR | ISSN (O) – 2582-6417
MARCH 2024 | Vol. 4 Issue 10
www.uijir.com

INTRODUCTION

The widespread use of antibiotics in pediatric populations has been a cornerstone of modern medicine, revolutionizing the treatment of bacterial infections in children and significantly reducing morbidity and mortality rates associated with such illnesses. However, alongside the benefits, concerns have emerged regarding the potential impact of this widespread antibiotic use on child immunity. Antibiotics are commonly prescribed to children for various conditions, including ear infections, strep throat, urinary tract infections, and respiratory tract infections. They work by killing or inhibiting the growth of bacteria, effectively treating bacterial infections. However, antibiotics are indiscriminate in their action, targeting both harmful bacteria causing infections and beneficial bacteria residing in the body, particularly in the gut.

One major concern surrounding the widespread use of antibiotics in pediatric populations is their potential to disrupt the delicate balance of the microbiota, which refers to the diverse community of microorganisms inhabiting the human body, especially the gut. The microbiota plays a crucial role in modulating the immune system, aiding in immune development, and maintaining immune homeostasis. Antibiotics can alter the composition and diversity of the microbiota, leading to dysbiosis, which is characterized by an imbalance in microbial populations. This dysbiosis may compromise immune function, making children more susceptible to infections and possibly contributing to the development of immune-related disorders.

Antibiotics can impact immune development and function beyond their effects on the microbiota. Research suggests that early-life exposure to antibiotics may interfere with the maturation of the immune system during critical developmental stages in infancy and childhood. This interference may impair the proper development of immune cells, such as T cells and B cells, and disrupt the intricate network of immune responses, potentially compromising the body's ability to defend against pathogens. The overuse and misuse of antibiotics can contribute to the emergence of antibiotic-resistant bacteria, posing a significant public health threat. Antibiotic resistance occurs when bacteria develop the ability to resist the effects of antibiotics, rendering these medications ineffective. This phenomenon not only complicates the treatment of bacterial infections in children but also undermines efforts to control infectious diseases globally.

While antibiotics have revolutionized the treatment of bacterial infections in pediatric populations, their widespread use raises legitimate concerns regarding their impact on child immunity. The disruption of the microbiota, interference with immune development, and the

DOI Link :: https://doi-ds.org/doilink/03.2024-51788784/UIJIR

© UIJIR | ISSN (O) – 2582-6417
MARCH 2024 | Vol. 4 Issue 10
www.uijir.com

emergence of antibiotic resistance are among the key concerns associated with antibiotic use in children. Therefore, it is essential for healthcare providers to judiciously prescribe antibiotics, considering both the potential benefits and risks, and to explore alternative strategies to manage infections while minimizing the impact on child immunity and overall health.

ANTIBIOTICS AND THE MICROBIOTA:

The composition and diversity of the microbiota, particularly in the gut, play a critical role in modulating immune homeostasis and function in children. Antibiotics, while effective in treating bacterial infections, can disrupt this delicate balance, leading to perturbations in the microbiota composition and diversity. Understanding how antibiotics alter the microbiota and the mechanisms through which these perturbations affect immune homeostasis and function is essential for comprehensively addressing the impact of antibiotics on child immunity.Antibiotics exert their effects by targeting and killing or inhibiting the growth of bacteria. However, they are broad-spectrum, meaning they can affect both pathogenic and commensal bacteria. In the context of the gut microbiota, antibiotics can lead to a reduction in the abundance and diversity of beneficial bacteria, such as Bacteroidetes and Firmicutes, while potentially allowing for the overgrowth of opportunistic pathogens. This disruption in microbial balance is commonly referred to as dysbiosis.

One mechanism through which antibiotics alter the composition and diversity of the microbiota is by directly killing susceptible bacteria. This can create ecological niches within the gut microbiota, allowing for the expansion of antibiotic-resistant strains or opportunistic pathogens that are resistant to the antibiotic being administered. Additionally, antibiotics can indirectly impact the microbiota by altering environmental conditions within the gut, such as pH levels and nutrient availability, which may further favor the growth of certain bacterial species over others.The perturbations in microbiota composition and diversity induced by antibiotics can have profound effects on immune homeostasis and function in children. The gut microbiota plays a crucial role in educating and shaping the developing immune system, particularly during infancy and early childhood. Commensal bacteria interact with immune cells in the gut-associated lymphoid tissue (GALT), influencing the maturation and function of various immune cell populations, including T cells, B cells, and regulatory T cells.

Disruption of the microbiota composition can lead to alterations in immune cell activation and cytokine production, potentially skewing immune responses towards inflammation or tolerance. For example, reductions in certain beneficial bacteria, such as those producing short-chain fatty

DOI Link :: https://doi-ds.org/doilink/03.2024-51788784/UIJIR

© UIJIR | ISSN (O) – 2582-6417
MARCH 2024 | Vol. 4 Issue 10
www.uijir.com

acids (SCFAs), can impair the development of regulatory T cells, which play a key role in immune tolerance and preventing autoimmune reactions. Dysbiosis may also compromise the integrity of the gut epithelial barrier, leading to increased permeability (leaky gut), which can further exacerbate immune dysregulation and inflammation.

Furthermore, alterations in the microbiota composition and diversity can impact the production of antimicrobial peptides and other molecules involved in host defense against pathogens. This may result in increased susceptibility to infections or the colonization of the gut by pathogenic bacteria, further disrupting immune homeostasis.

Antibiotics can significantly alter the composition and diversity of the microbiota in children, leading to dysbiosis and perturbations in immune homeostasis and function. Mechanisms underlying these effects include direct killing of susceptible bacteria, selection for antibiotic-resistant strains, and alterations in environmental conditions within the gut. Understanding these mechanisms is crucial for developing strategies to mitigate the adverse effects of antibiotics on child immunity while preserving their therapeutic benefits against bacterial infections. Additionally, efforts to restore and maintain a healthy gut microbiota following antibiotic treatment may hold promise for promoting immune health in children.

ANTIBIOTICS AND IMMUNE DYSREGULATION

The effects of antibiotics on immune cell populations, cytokine profiles, and immune signaling pathways in children are multifaceted and can have significant implications for immune function, susceptibility to infections, and the development of immune-related disorders. Understanding these effects requires an examination of how antibiotics impact various aspects of the immune system and how dysregulation induced by antibiotic exposure may contribute to adverse health outcomes in children. Antibiotics, while primarily targeting bacteria, can also influence immune cell populations in children. For instance, studies have shown that antibiotics can alter the abundance and distribution of immune cells, including T cells, B cells, dendritic cells, and macrophages. These changes may result from direct effects on immune cells or indirect effects mediated through alterations in the microbiota and gut environment. Antibiotic-induced shifts in immune cell populations can affect immune surveillance, response to pathogens, and the maintenance of immune homeostasis.

Antibiotics can modulate cytokine profiles, which are critical mediators of immune responses. Cytokines are signaling molecules secreted by immune cells that regulate inflammation, immune cell activation, and communication between different cell types. Antibiotics may

DOI Link :: https://doi-ds.org/doilink/03.2024-51788784/UIJIR

influence cytokine production and release, leading to alterations in the balance between pro-inflammatory and anti-inflammatory cytokines. Dysregulation of cytokine profiles can disrupt immune responses, impair tissue repair mechanisms, and contribute to chronic inflammation or immune-related disorders in children. Antibiotics can impact immune signaling pathways, which are intricate networks of molecular interactions that regulate immune cell activation, differentiation, and function. Antibiotics may interfere with these signaling pathways by modulating gene expression, protein synthesis, or post-translational modifications. For example, antibiotics may affect the activation of transcription factors such as nuclear factor kappa B (NF-κB) or signal transducer and activator of transcription (STAT) proteins, which play crucial roles in orchestrating immune responses. Dysregulation of immune signaling pathways can perturb immune cell function, impair host defense mechanisms, and increase susceptibility to infections or autoimmune reactions.

The dysregulation of immune cell populations, cytokine profiles, and immune signaling pathways induced by antibiotics may contribute to increased susceptibility to infections and the development of immune-related disorders in children. For instance, alterations in immune cell populations and cytokine profiles can weaken immune responses, making children more vulnerable to infections by opportunistic pathogens or pathogens resistant to antibiotics. Additionally, dysregulation of immune signaling pathways can promote chronic inflammation, tissue damage, and the development of autoimmune diseases or allergic conditions.The effects of antibiotics on immune cell populations, cytokine profiles, and immune signaling pathways in children are complex and multifaceted. Antibiotic-induced immune dysregulation can compromise immune function, increase susceptibility to infections, and contribute to the development of immune-related disorders. Understanding these effects is essential for optimizing antibiotic use, minimizing adverse health outcomes, and promoting immune health in children. Future research efforts should focus on elucidating the underlying mechanisms of antibiotic-induced immune dysregulation and identifying strategies to mitigate its impact on child immunity and overall well-being.

LONG-TERM CONSEQUENCES

A review of epidemiological studies and animal models investigating the long-term consequences of early-life antibiotic exposure on immune resilience and health outcomes in later life reveals a complex interplay between antibiotic use during critical developmental stages

DOI Link :: https://doi-ds.org/doilink/03.2024-51788784/UIJIR

© UIJIR | ISSN (O) – 2582-6417
MARCH 2024 | Vol. 4 Issue 10
www.uijir.com

and subsequent health outcomes. Epidemiological studies have provided evidence linking early-life antibiotic exposure to an increased risk of immune-related disorders, such as asthma, allergies, inflammatory bowel disease, and autoimmune diseases, later in life. Animal models have further elucidated potential underlying mechanisms and shed light on the long-term effects of early-life antibiotic exposure on immune resilience. Several epidemiological studies have reported associations between early-life antibiotic exposure and an elevated risk of immune-related disorders in children and adults. These associations suggest that antibiotic use during infancy and childhood may disrupt immune development and increase susceptibility to immune-mediated diseases later in life. Animal models have corroborated these findings, demonstrating that early-life antibiotic exposure can alter the composition and function of the gut microbiota, impair immune maturation, and predispose individuals to immune dysregulation and inflammatory conditions in adulthood.

The potential mechanisms underlying the observed associations between early-life antibiotic exposure and long-term health outcomes involve perturbations in the gut microbiota, immune dysregulation, and alterations in immune programming during critical developmental periods. Antibiotics disrupt the balance of the gut microbiota, leading to dysbiosis and alterations in microbial diversity and composition. This dysbiosis can have profound effects on immune function, including impaired immune cell development, dysregulated cytokine production, and compromised immune tolerance.

Early-life antibiotic exposure can disrupt immune programming during critical developmental stages, leading to long-lasting alterations in immune function and responsiveness. Antibiotics may interfere with the development of immune regulatory mechanisms, such as regulatory T cells, which play a crucial role in maintaining immune homeostasis and preventing autoimmunity. Dysregulation of immune programming during infancy and childhood can have enduring effects on immune resilience and increase susceptibility to immune-related disorders later in life. The implications of these findings for public health are significant, highlighting the importance of judicious antibiotic use in infancy and childhood to preserve immune health and minimize the risk of long-term adverse health outcomes. Strategies aimed at optimizing antibiotic prescribing practices, promoting alternatives to antibiotics when appropriate, and enhancing microbial stewardship in healthcare settings are essential for mitigating the impact of early-life antibiotic exposure on immune resilience and reducing the burden of immune-related diseases in the population.

DOI Link :: https://doi-ds.org/doilink/03.2024-51788784/UIJIR

© UIJIR | ISSN (O) – 2582-6417
MARCH 2024 | Vol. 4 Issue 10
www.uijir.com

STRATEGIES TO MITIGATE ADVERSE EFFECTS:

An overview of potential interventions and strategies to mitigate the adverse effects of antibiotics on child immunity involves a multifaceted approach aimed at preserving immune resilience while maintaining the therapeutic efficacy of antibiotics. Key considerations include the use of probiotics, prebiotics, and alternative approaches to promote a healthy microbiota and support immune function in children. Probiotics, which are live microorganisms that confer health benefits when consumed in adequate amounts, have been extensively studied for their potential to mitigate the adverse effects of antibiotics on the gut microbiota and immune system. By replenishing beneficial bacteria that may be depleted by antibiotics, probiotics can help restore microbial balance, enhance immune function, and reduce the risk of antibiotic-associated gastrointestinal symptoms, such as diarrhea. Clinical trials have shown that certain probiotic strains, such as Lactobacillus and Bifidobacterium species, can effectively prevent antibiotic-associated diarrhea and promote gut health in children receiving antibiotics.

In addition to probiotics, prebiotics, which are non-digestible dietary fibers that serve as a fuel source for beneficial bacteria in the gut, can also support the restoration of the microbiota following antibiotic treatment. By promoting the growth of beneficial bacteria, prebiotics can help reestablish microbial diversity and enhance immune function. Incorporating prebiotic-rich foods, such as fruits, vegetables, whole grains, and legumes, into children's diets can provide a natural source of prebiotics to support gut health and immune resilience. Alternative approaches to antibiotics, such as antimicrobial stewardship practices, targeted antimicrobial therapies, and the use of immune-modulating agents, can help minimize the need for antibiotic treatment and reduce the risk of antibiotic-associated complications in children. Antimicrobial stewardship programs aim to optimize antibiotic use by promoting appropriate prescribing practices, minimizing unnecessary antibiotic exposure, and preventing the emergence of antibiotic resistance. Targeted antimicrobial therapies, such as narrow-spectrum antibiotics or combination therapies, can be used to selectively target pathogenic bacteria while minimizing disruption to the commensal microbiota. Additionally, immune-modulating agents, such as cytokine inhibitors or immunomodulatory drugs, may offer alternative approaches to managing infections and supporting immune function in children with specific immune-related disorders.

Optimizing antibiotic use and implementing strategies to support immune resilience in children are essential for mitigating the adverse effects of antibiotics on child immunity. By incorporating probiotics, prebiotics, and alternative approaches into clinical practice, healthcare providers can help preserve immune health, minimize the risk of antibiotic-associated

DOI Link :: https://doi-ds.org/doilink/03.2024-51788784/UIJIR

complications, and promote long-term well-being in pediatric populations. Continued research efforts are needed to further elucidate the effectiveness and safety of these interventions and their impact on immune resilience and health outcomes in children.

CONCLUSION

Key findings from research on the interactions between antibiotics, child immunity, and long-term health outcomes highlight the importance of judicious antibiotic use, promoting immune resilience, and implementing strategies to mitigate the adverse effects of antibiotics on child health. In clinical practice, healthcare providers should prioritize antibiotic prescribing practices that minimize unnecessary antibiotic exposure, avoid broad-spectrum antibiotics when possible, and consider the potential impact of antibiotics on the developing immune system. Probiotics, prebiotics, and alternative approaches can be incorporated into treatment plans to support immune resilience and mitigate antibiotic-associated complications in children. Public health policies should focus on promoting antimicrobial stewardship initiatives that educate healthcare providers, patients, and caregivers about the appropriate use of antibiotics, encourage the development of guidelines for antibiotic prescribing in pediatric populations, and promote research into alternative treatment modalities that minimize the use of antibiotics.

Antibiotic stewardship initiatives should prioritize efforts to reduce antibiotic overuse and misuse, promote antimicrobial stewardship best practices in healthcare settings, and support the development of novel diagnostic tools and treatment strategies that minimize the need for antibiotics. By optimizing antibiotic use and supporting immune resilience in children, these initiatives can help preserve the effectiveness of antibiotics, reduce the risk of antibiotic resistance, and promote better long-term health outcomes. Further research is needed to elucidate the complex interactions between antibiotics, child immunity, and long-term health outcomes. Future studies should focus on understanding the mechanisms underlying the effects of antibiotics on immune development and function in children, identifying biomarkers of immune resilience and susceptibility to antibiotic-associated complications, and evaluating the effectiveness and safety of interventions aimed at mitigating the adverse effects of antibiotics on child health.

In conclusion, research on antibiotics, child immunity, and long-term health outcomes underscores the importance of optimizing antibiotic use, promoting immune resilience, and implementing strategies to mitigate the adverse effects of antibiotics on child health. By prioritizing these efforts in clinical practice, public health policies, and antibiotic stewardship

DOI Link :: https://doi-ds.org/doilink/03.2024-51788784/UIJIR

© UIJIR | ISSN (O) – 2582-6417
MARCH 2024 | Vol. 4 Issue 10
www.uijir.com

initiatives, healthcare providers and policymakers can help preserve the effectiveness of antibiotics and promote better long-term health outcomes for children.

DOI Link :: https://doi-ds.org/doilink/03.2024-51788784/UIJIR

© UIJIR | ISSN (O) – 2582-6417
MARCH 2024 | Vol. 4 Issue 10
www.uijir.com

REFERENCES

1. Alexander VN, Northrup V, Bizzarro MJ. Antibiotic exposure in the newborn intensive care unit and the risk of necrotizing enterocolitis. *J Pediatr-Us.* (2011) 159:392–7. 10.1016/j.jpeds.2011.02.035 [
2. Arpaia N, Campbell C, Fan XY, Dikiy S, van der Veeken J, deRoos P, et al.. Metabolites produced by commensal bacteria promote peripheral regulatory T-cell ge*neration*. *Nature*. (2013) 504:451–5. 10.1038/nature12726
3. Clarke TB, Davis KM, Lysenko ES, Zhou AY, Yu YM, Weiser JN. Recognition of peptidoglycan from the microbiota by Nod1 enhances systemic innate immunity. *Nat Med.* (2010) 16:228–U137. 10.1038/nm.2087
4. Cotten CM, Taylor S, Stoll B, Goldberg RN, Hansen NI, Sanchez PJ, et al.. Prolonged duration of initial empirical antibiotic treatment is associated with increased rates of necrotizing enterocolitis and death for extremely low birth weight infants. *Pediatrics.* (2009) 123:58–66. 10.1542/peds.2007-3423
5. De Kleer, Willems F, Lambrecht B, Goriely S. Ontogeny of myeloid cells. *Front Immunol.* (2014) 5:423. 10.3389/fimmu.2014.00423
6. Deshmukh HS, Liu YH, Menkiti OR, Mei JJ, Dai N, O'Leary CE, et al.. The microbiota regulates neutrophil homeostasis and host resistance to Escherichia coli K1 sepsis in neonatal mice. *Nat Med.* (2014) 20:528–34. 10.1038/nm.3542
7. Edelson BT, Kc W, Juang R, Kohyama M, Benoit LA, Klekotka PA, et al.. Peripheral CD103+ dendritic cells form a unified subset developmentally related to CD8alpha+ conventional dendritic cells. *J Exp Med.* (2010) 207:823–36. 10.1084/jem.20091627
8. Fouhse JM, Yang KY, More-Bayona J, Gao YH, Goruk S, Plastow G, et al.. Neonatal exposure to amoxicillin alters long-term immune response despite transient effects on gut-microbiota in piglets. *Front Immunol.* (2019) 10:2059. 10.3389/fimmu.2019.02059
9. Francino MP. Early development of the gut microbiota and immune health. *Pathogens*. (2014) 3:769–90. 10.3390/pathogens3030769 [
10. Gonzalez-Perez G, Hicks AL, Tekieli TM, Radens CM, Williams BL, Lamouse-Smith ESN. Maternal antibiotic treatment impacts development of the neonatal intestinal microbiome and antiviral immunity. *J Immunol.* (2016) 196:3768–79. 10.4049/jimmunol.1502322

DOI Link :: https://doi-ds.org/doilink/03.2024-51788784/UIJIR

© UIJIR | ISSN (O) – 2582-6417
MARCH 2024 | Vol. 4 Issue 10
www.uijir.com

11. Gonzalez-Perez G, Lamouse-Smith ESN. Gastrointestinal microbiome dysbiosis in infant mice alters peripheral CD8(+) T cell receptor signaling. *Front Immunol.* (2017) 8:265. 10.3389/fimmu.2017.
12. Gopinath S, Kim MV, Rakib T, Wong PW, van Zandt M, Barry NA, et al.. Topical application of aminoglycoside antibiotics enhances host resistance to viral infections in a microbiota-independent manner. *Nat Microbiol.* (2018) 3:611–21. 10.1038/s41564-018-0138-2
13. Gray J, Oehrle K, Worthen G, Alenghat T, Whitsett J, Deshmukh H. Intestinal commensal bacteria mediate lung mucosal immunity and promote resistance of newborn mice to infection. *Sci Transl Med.* (2017) 9:eaaf9412. 10.1126/scitranslmed.aaf9412
14. Han D, Walsh MC, Kim KS, Hong SW, Lee J, Yi J. Microbiota-independent ameliorative effects of antibiotics on spontaneous Th2-associated pathology of the small intestine. *PLoS ONE.* (2015) 10:e0118795. 10.1371/journal.pone.0118795
15. Ichinohe T, Pang IK, Kumamoto Y, Peaper DR, Ho JH, Murray TS, et al.. Microbiota regulates immune defense against respiratory tract influenza A virus infection. *Proc Natl Acad Sci USA.* (2011) 108:5354–9. 10.1073/pnas.1019378108
16. Jernberg C, Lofmark S, Edlund C, Jansson JK. Long-term impacts of antibiotic exposure on the human intestinal microbiota. *Microbiology.* (2010) 156:3216–23. 10.1099/mic.0.040618-0
17. Khan R, Petersen FC, Shekhar S. Commensal bacteria: an emerging player in defense against respiratory pathogens. *Front Immunol.* (2019) 10:1203. 10.3389/fimmu.2019.01203
18. Korpela K, Salonen A, Virta LJ, Kekkonen RA, Forslund K, Bork P, et al.. Intestinal microbiome is related to lifetime antibiotic use in Finnish pre-school children. *Nat Commun.* (2016) 7:10410. 10.1038/ncomms10410
19. Kuppala VS, Meinzen-Derr J, Morrow AL, Schibler KR. Prolonged initial empirical antibiotic treatment is associated with adverse outcomes in premature infants. *J Pediatr.* (2011) 159:720–5. 10.1016/j.jpeds.2011.05.033
20. Lamouse-Smith ES, Tzeng A, Starnbach MN. The intestinal flora is required to support antibody responses to systemic immunization in infant and germ free mice. *PLoS ONE.* (2011) 6:e27662. 10.1371/journal.pone.0027662

DOI Link :: https://doi-ds.org/doilink/03.2024-51788784/UIJIR

21. Lynn MA, Tumes DJ, Choo JM, Sribnaia A, Blake SJ, Leong LEX, et al.. Early-life antibiotic-driven dysbiosis leads to dysregulated vaccine immune responses in mice. *Cell Host Microbe.* (2018) 23:653–60. 10.1016/j.chom.2018.04.009
22. Man WH, Clerc M. W., Piters AAD, van Houten MA, M.Chu LJN, Kool J, et al.. Loss of microbial topography between oral and nasopharyngeal microbiota and development of respiratory infections early in life. *Am J Resp Crit Care.* (2019) 200:760–70. 10.1164/rccm.201810-1993OC
23. Miller JE, Wu CS, Pedersen LH, de Klerk N, Olsen J, Burgner DP. Maternal antibiotic exposure during pregnancy and hospitalization with infection in offspring: a population-based cohort study. *Int J Epidemiol.* (2018) 47:561–71. 10.1093/ije/dyx272
24. Nair AB, Jacob S. A simple practice guide for dose conversion between animals and human. *J Basic Clin Pharm.* (2016) 7:27–31. 10.4103/0976-0105.177703
25. Olin A, Henckel E, Chen Y, Lakshmikanth T, Pou C, Mikes J, et al.. Stereotypic Immune system development in newborn children. *Cell.* (2018) 174:1277–92.e14. 10.1016/j.cell.2018.06.045
26. Oosterlo BC, van't Land B, de Jager W, Rutten NB, Klopping M, Garssen J. V, et al.. Neonatal antibiotic treatment is associated with an altered circulating immune marker profile at 1 year of age. *Front Immunol.* (2020) 10:2939. 10.3389/fimmu.2019.02939
27. Patrick DM, Sbihi H, Dai DLY, Al Mamun A, Rasali D, Rose C, et al.. Decreasing antibiotic use, the gut microbiota, and asthma incidence in children: evidence from population-based and prospective cohort studies. *Lancet Respir Med.* (2020) 2600:30052–7. 10.1016/S2213-2600(20)30052-7
28. Rogawski ET, Westreich D, Becker-Dreps S, Adair LS, Sandler RS, Sarkar R, et al.. The effect of early life antibiotic exposures on diarrheal rates among young children in Vellore, India. *Pediatr Infect Dis J.* (2015) 34:583–8. 10.1097/INF.0000000000000679
29. Roubaud-Baudron C, Ruiz VE, Swan AM, Vallance BA, Ozkul C, Pei ZH, et al.. Long-term effects of early-life antibiotic exposure on resistance to subsequent bacterial infection. *Mbio.* (2019) 10:e02820–19. 10.1128/mBio.02820-
30. Torres D, Munoz T, Bancalari A, Manriquez C. Prolonged initial empirical antibiotic treatment and the risk of morbidity and mortality in very low birthweight infants. *Rev Chil Pediatr-Chi.* (2018) 89:600–5. 10.4067/S0370-41062018005000807

DOI Link :: https://doi-ds.org/doilink/03.2024-51788784/UIJIR

© UIJIR | ISSN (O) – 2582-6417
MARCH 2024 | Vol. 4 Issue 10
www.uijir.com

Zhu DP, Xiao S, Yu JL, Ai Q, He Y, Cheng C, et al.. Effects of one-week empirical antibiotic therapy on the early development of gut microbiota and metabolites in preterm infants. *Sci Rep.* (2017) 7:8025. 10.1038

DOI Link :: https://doi-ds.org/doilink/03.2024-51788784/UIJIR